

NEUROSURGICAL ASSOCIATES, L.L.C  
NAM YOON M.D., P.C.

5171 South Cottonwood Street, Suite 950  
Murray, Utah 84157  
Phone (801) 507-9555  
Toll Free (866) 804-2193  
Fax (801) 507-9550

**DISCHARGE INSTRUCTIONS FOLLOWING A LUMBAR MICRODISCECTOMY**

You had a surgical procedure call a lumbar microdiscectomy. During the procedure Dr. Yoon made a small incision(s) on your low back. Then the disc material that was pushing on your spinal nerve(s) causing your pain or leg weakness was removed. The rest of the disc is left in place. These instructions provide additional information regarding your care at home.

**Incision Care:**

- Starting two days after surgery, you may shower with mild soap and shampoo daily. Gently wash your incision and pat dry. This is the only time you may touch your incision.
- Your incision was closed with absorbable sutures. Steri-strips were placed over the skin. The steri-strips should be removed after 1-2 weeks if they are still in place.
- Do not apply ointments, lotions or creams to your incision.
- Apply an ice pack or a clean bag of frozen peas to your incision every 30 minutes to help reduce the swelling and discomfort, as needed.
- Do not use a hot tub, go swimming or take a bath until your incisions are completely healed (about 4 weeks for patients).
- Stop smoking as this delays healing and may cause a wound infection.

**Activities:**

- Start with light activity around the house for the first 2 days you are home.
- Gradually increase your activity starting with a short walk 1-2 times a day.
- Allow your body time to heal by resting for short periods during the day.
- Avoid contact sports, skating, bike riding or other activities for 6 weeks.
- You may **not** drive when taking pain medications and until cleared by Dr. Yoon.
- Avoid lifting, pushing or pulling heavy objects (more than 10 lbs) for 6-12 weeks.
- Avoid bending over or twisting to pick up things.
- Avoid sitting in soft chairs or slumping while you are sitting.
- Be sure to get up and move around/stretch every 30 minutes while sitting.

**Nutrition:**

- Eat plenty of fruits and vegetables to prevent constipation.

**Medications:**

- Take your medications as prescribed and gradually decrease pain medications as your pain improves.
- You may need to take a stool softener (Colace) or a laxative that you can buy at a pharmacy until your bowels return to normal.
- You may need to use a suppository (Dulcolax or glycerin) or an enema if you have not had a bowel movement in 3 days.
- **Notes about pain medications and opioids (Norco, hydrocodone, oxycodone, Percocet, Dilaudid):** These medications can be habit forming and addicting, and can lead to overdose and possibly death. Do not combine these medications with alcohol, and if you take other sedating medications (muscle relaxers, antianxiety meds, for example) or have sleep apnea, take extra care while using these medicines. We will start decreasing your dose after a week or so, and will only be able to prescribe pain medications for a maximum of three months after surgery. After that time, you will need to contact your prior provider for assistance.

**Follow Up:**

- Unless this has already been arranged, call Dr. Yoon's office when you get home to schedule your follow-up appointment.
- Follow up with your Primary Care Physician for all medical issues.

**Call your doctor or return to the emergency room if you experience any of the following:**

- Constipation- no bowel movement for more than 3 days.
- Difficulty moving or weakness of your legs.
- Back pain that is not well controlled on your pain medications.
- **A fever above 100F.**
- Redness, swelling, odor or drainage at your incision site.
- Loss of bowel or bladder control.
- Headaches in an upright position which resolve with lying down.
- Difficulty feeling your legs.
- Difficulty walking.

